arXiv:0709.0785v2 [math.AT] 25 Jul 2008

A DESCRIPTION BASED ON SCHUBERT CLASSES OF COHOMOLOGY OF FLAG MANIFOLDS

MASAKI NAKAGAWA*

Abstract. We describe the integral cohomology rings of the flag manifolds of types B_n, D_n, G_2 and F_4 in terms of their Schubert classes. The main tool is the divided difference operators of Bernstein-Gelfand-Gelfand and Demazure. As an application, we compute the Chow rings of the corresponding complex algebraic groups, recovering thereby the results of R. Marlin.

1. Introduction

Let K be a compact connected Lie group and T its maximal torus. The homogeneous space K/T, called the flag manifold, plays an important role in algebraic topology, algebraic geometry and representation theory.

In this paper, we are concerned with the integral cohomology of the flag manifold K/T. As is well known, there are two descriptions of the integral cohomology of flag manifold. The first one is the "*Borel presentation*," due to A. Borel [2], which identifies the rational cohomology ring of K/T with the quotient ring of a polynomial ring by its ideal generated by W-invariants of positive degrees, where W is the Weyl group of K. Combining Borel's result and the known structures of the mod p cohomology rings of K, H. Toda gave general descriptions of the integral cohomology rings of K/T for all K simple [18]. So far the integral cohomology rings of flag manifolds for all compact simply connected simple Lie groups are determined (see [2], [3], [19], [14], [15]). The second is the "*Schubert presentation*" which describes the integral cohomology $H^*(K/T;\mathbb{Z})$ in terms of the Schubert classes corresponding to the Schubert varieties derived from the Bruhat decomposition of $G = K^{\mathbb{C}}$, the complexification of K.

2000 *Mathematics Subject Classification.* Primary 57T15; Secondary 14M15.

Key words and phrases. flag manifolds, Schubert calculus, Chow rings.

* Partially supported by the Grant-in-Aid for Scientific Research (C) 18540106, Japan Society for the Promotion of Science.

In the Borel presentation, which is given by generators and relations, the ring structure of $H^*(K/T;\mathbb{Z})$ can be relatively easy to obtain. However, the generators in this presentation have little geometric meaning. In contrast, in the Schubert presentation, the Schubert classes correspond to the geometric objects - the Schubert varieties - and they form an additive basis for $H^*(K/T;\mathbb{Z})$ ([5]). As a disadvantage, the multiplicative structure among them is closely related to the intersection multiplicities, and is highly complicated in general.

Up to now, there have been several attempts to establish a connection between the two descriptions for some types of spaces (see, e.g., [1], [8], [11]). The main aim of this paper also falls in this category. More precisely, we express the ring generators in the Borel presentation of $H^*(K/T;\mathbb{Z})$ for $K = SO(n), G_2$ and F_4 in terms of Schubert classes. For this, we make use of the "*divided difference operators*" introduced independently by Bernstein-Gelfand-Gelfand [1] and Demazure [6]. For K as above, there exist extra generators of degrees greater than two. So we cannot apply the divided difference operators directly to these higher generators. Fortunately, using the classical fact that, rationally, the cohomology of K/T is generated as a ring by two dimensional elements and the integral cohomology of K/T has no torsion, we can carry out the computation. An additional aim of this paper is to apply our results to recovering the Chow rings of the complex algebraic groups $\mathrm{SO}(n), \mathrm{Spin}(n), \mathrm{G}_2$ and F_4, which were originally computed by R. Marlin [13]. (In this paper, we denote the compact Lie groups, e.g., by $SO(n), Spin(n), G_2, F_4$, while their complexifications by $\mathrm{SO}(n), \mathrm{Spin}(n), \mathrm{G}_2, \mathrm{F}_4$ respectively.) In order to determine the Chow rings of the corresponding flag manifolds, Marlin relied on the result of Demazure [6] which describes them as the "*cohomology rings of the root system*," and he made elaborate computations. In this paper, we simplify Marlin's computations, using the Borel presentation of $H^*(K/T;\mathbb{Z})$ and our result mentioned above.

The paper is organized as follows. In Section 2, we briefly review the cohomology of flag manifolds, emphasizing the difference between the Borel presentation and the Schubert presentation. In Section 3, we introduce the divided difference operators of Bernstein-Gelfand-Gelfand and Demazure and collect the results used later. Sections 4 to 6 are devoted to computations, and we obtain there the main results of this paper (see Propositions 4.3, 4.7, 5.3 and 6.5). In Section 7, following Grothendieck's remark ([9], p.21, REMARQUES 2°), we compute the Chow rings of $\mathrm{SO}(n), \mathrm{Spin}(n), \mathrm{G}_2$ and F_4 (see Theorems 7.2, 7.3, 7.4 and 7.1).

We observe that the method of this paper can also be applied to the exceptional Lie groups E_6, E_7 and E_8. Indeed, we succeeded in computing the Chow rings of the complex algebraic groups E_6 and E_7 in [12].

Acknowledgments. Firstly, we thank Július Korbaš for reading carefully the manuscript and giving us useful comments and William Fulton for kindly pointing out an error concerning the Schubert classes in the earlier version of the manuscript. Secondly, we thank Mamoru Mimura for giving us various suggestions. Finally, we thank Shizuo Kaji for the program using Maple, which confirmed our computations.

2. THE COHOMOLOGY OF FLAG MANIFOLDS

In this section we briefly review the Borel presentation and the Schubert presentation of the cohomology of flag manifolds.

We introduce the notation that is needed in the sequel.

K: a compact simply connected simple Lie group of rank l;
T: a maximal torus of K;
$G = K^{\mathbb{C}}$: a complexification of K;
B: a Borel subgroup containing T;
$\mathfrak{t}$: the Lie algebra of T, $\mathfrak{t}^*$: the dual space of $\mathfrak{t}$;
$(\,\cdot\,|\,\cdot\,)$: the invariant inner product on $\mathfrak{t}$ (or on $\mathfrak{t}^*$);
$\Delta \subset \mathfrak{t}^*$: the root system with respect to T;
Δ^+: a set of positive roots, $\Delta^- = -\Delta^+$;
$\Pi = \{\alpha_1, \ldots, \alpha_l\}$: the system of simple roots;
$\alpha^\vee = \dfrac{2\alpha}{(\alpha|\alpha)}$: the coroot corresponding to $\alpha \in \Delta$;
ω_i $(1 \le i \le l)$: the ith fundamental weight, satisfying $(\omega_i|\alpha_j^\vee) = \delta_{ij}$;
$s_i = s_{\alpha_i}$ $(1 \le i \le l)$: the reflection corresponding to the simple root α_i;
$W = W(K)$: the Weyl group of K generated by the simple reflections $S = \{s_1, \ldots, s_l\}$;
$l(w)$: the length of an element $w \in W$ with respect to $\{s_1, \ldots, s_l\}$;
w_0: the longest element of W;
$e_i(x_1, \ldots, x_n)$: the ith elementary symmetric function in variables $x_1, \ldots, x_n$.

Now we review the Borel presentation. The inclusion $T \hookrightarrow K$ induces the classical fibration

$$K/T \xrightarrow{\iota} BT \xrightarrow{\rho} BK,$$

where BT (resp. BK) denotes the classifying space of T (resp. K). The induced homomorphism

$$c = \iota^* : H^*(BT;\mathbb{Z}) \longrightarrow H^*(K/T;\mathbb{Z}) \tag{2.1}$$

is called the *characteristic homomorphism* and plays a crucial role in Borel's work. The Weyl group W acts naturally on T, hence on $H^2(BT;\mathbb{Z})$. We extend this natural action of W to the whole $H^*(BT;\mathbb{Z})$ and also to $H^*(BT;\mathbb{F}) = H^*(BT;\mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{F}$, where $\mathbb{F}$ is any field. We denote by $H^*(BT;\mathbb{Z})^W$ (resp. $H^*(BT;\mathbb{F})^W$) the ring of W-invariants in $H^*(BT;\mathbb{Z})$ (resp. $H^*(BT;\mathbb{F})$). Then one of the main results of Borel can be stated as follows.

Theorem 2.1 (Borel [2]). *Let $\mathbb{F}$ be a field of characteristic zero. Then the characteristic homomorphism induces an isomorphism*

$$\bar{c} : H^*(BT;\mathbb{F})/(H^+(BT;\mathbb{F})^W) \longrightarrow H^*(K/T;\mathbb{F}),$$

where $(H^+(BT;\mathbb{F})^W)$ is the ideal in $H^*(BT;\mathbb{F})$ generated by the W-invariants of positive degrees.

In particular, one can reduce the computation of the rational cohomology ring $H^*(K/T;\mathbb{Q})$ to that of the ring of invariants $H^*(BT;\mathbb{Q})^W$. Observe that $H^*(K/T;\mathbb{Q})$ is generated by $H^2(K/T;\mathbb{Q})$ as a ring. In order to determine the integral cohomology ring $H^*(K/T;\mathbb{Z})$, we need further considerations. In [18], Toda established a method to describe the integral cohomology ring $H^*(K/T;\mathbb{Z})$ by a minimal system of generators and relations, from the mod p cohomology rings $H^*(K;\mathbb{Z}/p\mathbb{Z})$ and the rational cohomology ring $H^*(K/T;\mathbb{Q})$. In general, besides the two-dimensional generators, there are extra generators of higher degrees, and hence the characteristic homomorphism c is not surjective over $\mathbb{Z}$ in that case. Along the lines of Toda's method, the integral cohomology rings of flag manifolds for all compact simply connected simple Lie groups have been computed (see [2], [3], [19], [14], [15]). However, as mentioned in the introduction, the generators have less geometric meaning in the Borel presentation.

We pass to reviewing the Schubert presentation. Recall the Bruhat decomposition,

$$G = \coprod_{w \in W} B\dot{w}B,$$

where $\dot{w}$ denotes any representative of w in $W = N_K(T)/T$, $N_K(T)$ is the normalizer of T in K. It induces a cell decomposition,

$$G/B = \coprod_{w \in W} B\dot{w}B/B,$$

where

$$X_w^\circ = B\dot{w}B/B \cong \mathbb{C}^{l(w)}$$

is called the *Schubert cell.* Note that we have a homeomorphism $K/T \cong G/B$ by the Iwasawa decomposition. The *Schubert variety* X_w is defined to be the closure of X_w°. Then it is known that

$$X_w = \coprod_{v \le w} X_v^\circ,$$

where $\le$ is the Bruhat-Chevalley ordering. The fundamental class $[X_w]$ of X_w lies in $H_{2l(w)}(G/B;\mathbb{Z})$. We define the cohomology class $Z_w \in H^{2l(w)}(G/B;\mathbb{Z})$ as the Poincaré dual of $[X_{w_0 w}] \in H_{2N-2l(w)}(G/B;\mathbb{Z})$, where N is the complex dimension of the flag manifold G/B. We call Z_w the *Schubert class.* The Schubert classes $\{Z_w\}_{w\in W}$ form an additive basis for $H^*(G/B;\mathbb{Z})$. We refer to $\{Z_w\}_{w\in W}$ as the *Schubert basis.* In order to complete the description of $H^*(G/B;\mathbb{Z})$, we have to compute the intersection multiplicities. Namely, given $u, v \in W$, we can put

$$Z_u \cdot Z_v = \sum_{\substack{w \in W, \\ l(u)+l(v)=l(w)}} a_{u,v}^w Z_w$$

for some integers $a_{u,v}^w$, and we have to determine these "structure constants" $a_{u,v}^w$. As for this problem, several results are available. For example, we have the following

Theorem 2.2 (Chevalley formula [5])**.** *If* $w \in W,\ \alpha \in \Pi$*, then we have the formula*

$$Z_{s_\alpha} \cdot Z_w = \sum_{\substack{\beta \in \Delta^+, \\ l(ws_\beta)=l(w)+1}} (\beta^\vee | \omega_\alpha) Z_{ws_\beta}.$$

For recent developments in the Schubert calculus, in particular, on multiplying Schubert classes, see also [7], [16].

3. Schubert calculus on flag manifolds

As reviewed in the previous section, there are two different ways of describing the integral cohomology ring of K/T, and therefore we have two bases for $H^*(K/T;\mathbb{Z})$ correspondingly. One is the "algebraic basis" derived from the Borel presentation and the other is the "geometric basis" $\{Z_w\}_{w\in W}$ consisting of the Schubert classes. It is interesting to know how these two bases are related. More precisely, we wish to express the ring generators obtained in the Borel presentation in terms of the Schubert basis. Our main tool will be the divided difference operators introduced independently by Bernstein-Gelfand-Gelfand [1] and Demazure [6]. We now recall their definition. For $\alpha \in \Delta$, we define an endomorphism of $H^*(BT;\mathbb{Z})$ by

$$\Delta_\alpha(u) = \frac{u - s_\alpha(u)}{\alpha}, \quad u \in H^*(BT;\mathbb{Z}).$$

Definition 3.1. *For $w \in W$, we define the operator*

$$\Delta_w = \Delta_{\alpha_{i_1}} \circ \cdots \circ \Delta_{\alpha_{i_k}}$$

on $H^*(BT;\mathbb{Z})$ lowering the degree by $2l(w)$, where $w = s_{i_1} \cdots s_{i_k}$ is a reduced decomposition of w.

One can show that the operator Δ_w is well defined, i.e., it is independent of the choice of the reduced decomposition of w.

Note that the divided difference operators $\Delta_\alpha, \alpha \in \Delta$, are characterized by the following two properties:

$$\Delta_\alpha(\omega_\beta) = \delta_{\alpha\beta}, \tag{3.1}$$

$$\Delta_\alpha(uv) = \Delta_\alpha(u)v + s_\alpha(u)\Delta_\alpha(v) \tag{3.2}$$

for $u, v \in H^*(BT;\mathbb{Z})$.

The characteristic homomorphism

$$c : H^*(BT;\mathbb{Z}) \longrightarrow H^*(K/T;\mathbb{Z})$$

can be described by the divided difference operators. Since $\{Z_w\}_{w\in W}$ is a $\mathbb{Z}$-basis for $H^*(K/T;\mathbb{Z})$, we can put

$$c(f) = \sum_{l(w)=k} a_w Z_w, \quad a_w \in \mathbb{Z}$$

for a polynomial $f \in H^{2k}(BT;\mathbb{Z})$. So we have to determine the coefficients a_w. This problem was solved independently by Bernstein-Gelfand-Gelfand [1] and Demazure [6].

Theorem 3.2 (Bernstein-Gelfand-Gelfand [1], Demazure [6])**.** *For a polynomial $f \in H^{2k}(BT;\mathbb{Z})$, we have*

$$c(f) = \sum_{l(w)=k} \Delta_w(f) Z_w.$$

In particular, for $\alpha \in \Pi$, we have

$$c(\omega_\alpha) = Z_{s_\alpha}.$$

In addition to this, using the divided difference operators, we can express an arbitrary Schubert class Z_w as a polynomial in the variables

Z_{s_i}. This is the Giambelli formula which we now recall (for details, see [10], Section 3). Consider the element

$$d = \prod_{\alpha \in \Delta^+} \alpha.$$

Then we have

Theorem 3.3 (Giambelli formula)**.** *The Schubert class* Z_w *corresponding to* $w \in W$ *is given by*

$$Z_w = c\left(\Delta_{w^{-1}w_0}\left(\frac{d}{|W|}\right)\right).$$

In Sections 4, 5 and 6, we exploit the above theorems to find the correspondence between "algebraic bases" and "geometric bases" in the cases of $K = SO(n), G_2$ and F_4.

4. THE CASES OF B_n AND D_n

In this section, we consider the special orthogonal group $SO(n)$. First we deal with the odd special orthogonal group $SO(2n+1)$. Let T^n be the standard maximal torus of $SO(2n+1)$,

$$T^n = \left\{ \begin{pmatrix} \cos 2\pi t_1 & -\sin 2\pi t_1 & & & & \\ \sin 2\pi t_1 & \cos 2\pi t_1 & & & & \\ & & \ddots & & & \\ & & & \cos 2\pi t_n & -\sin 2\pi t_n & \\ & & & \sin 2\pi t_n & \cos 2\pi t_n & \\ & & & & & 1 \end{pmatrix} \right\}.$$

Then we have an isomorphism:

$$H^*(BT^n; \mathbb{Z}) = \mathbb{Z}[t_1, t_2, \ldots, t_n].$$

The system of simple roots is given by

$$\Pi = \{\alpha_1 = t_1 - t_2, \alpha_2 = t_2 - t_3, \ldots, \alpha_{n-1} = t_{n-1} - t_n, \alpha_n = t_n\}.$$

The corresponding fundamental weights $\{\omega_i\}_{1 \le i \le n}$ are given by

$$\begin{aligned} \omega_i &= t_1 + t_2 + \cdots + t_i \quad (1 \le i \le n-1), \\ \omega_n &= \frac{1}{2}(t_1 + \cdots + t_n). \end{aligned} \tag{4.1}$$

Let $s_i\,(1 \le i \le n)$ be the reflection corresponding to the simple root $\alpha_i\,(1 \le i \le n)$. Then the Weyl group $W(SO(2n+1))$ is finite and is generated by $s_i\,(1 \le i \le n)$ which act on $\{t_i\}_{1\le i\le n}$ as permutations and signs changes:

$$\begin{aligned} W(SO(2n+1)) &= \langle s_1, s_2, \ldots, s_n\rangle \\ &\cong \mathcal{S}_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n, \end{aligned} \tag{4.2}$$

where $\mathcal{S}_n$ is the symmetric group of n letters and $\ltimes$ means the semi-direct product.

Now we recall the Borel presentation of $H^*(SO(2n+1)/T^n;\mathbb{Z})$ that had been probably known, in some form, already to Borel. However, in an explicit form, it first appeared in [19] as far as the author knows.

Theorem 4.1 (Toda-Watanabe [19], Theorem 2.1)**.** *The integral cohomology ring of $SO(2n+1)/T^n$ is*

$$H^*(SO(2n+1)/T^n;\mathbb{Z}) = \mathbb{Z}[t_1, t_2, \ldots, t_n, \gamma_1, \gamma_2, \ldots, \gamma_n] \Big/ \left(\begin{array}{l} c_i - 2\gamma_i \ (1 \le i \le n), \\ \gamma_{2k} + \displaystyle\sum_{i=1}^{2k-1} (-1)^i \gamma_i \gamma_{2k-i} \ (1 \le k \le n) \end{array} \right),$$

where we denote by the same symbols $t_i \in H^2(SO(2n+1)/T^n;\mathbb{Z})$ the images of $t_i \in H^2(BT^n;\mathbb{Z})$ under the homomorphism c, $c_i = e_i(t_1, \ldots, t_n)$ $(1 \le i \le n)$, and $\gamma_i = 0$ for $i > n$.

We wish to express the algebra generators $\{t_1, \ldots, t_n, \gamma_1, \ldots, \gamma_n\}$ in terms of Schubert classes. For simplicity, we denote the Schubert class corresponding to the element $w = s_{i_1}s_{i_2}\cdots s_{i_k}$ by $Z_{i_1 i_2 \cdots i_k}$, although the reduced decomposition of a Weyl group element may not be unique. The correspondence between elements of degree 2 is easy. By (4.1), we have

$$\begin{aligned} t_1 &= \omega_1, \\ t_i &= -\omega_{i-1} + \omega_i \quad (2 \le i \le n-1), \\ t_n &= -\omega_{n-1} + 2\omega_n. \end{aligned} \tag{4.3}$$

Since $c(\omega_i) = Z_i$, it follows from (4.3) that

$$\begin{aligned} t_1 &= Z_1, \\ t_i &= -Z_{i-1} + Z_i \quad (2 \le i \le n-1), \\ t_n &= -Z_{n-1} + 2Z_n. \end{aligned} \tag{4.4}$$

Next, for γ_k $(1 \le k \le n)$, we can put

$$\gamma_k = \sum_{l(w)=k} a_w Z_w$$

for some integers a_w. We need to determine the coefficients a_w. To this end, we make use of the divided difference operators recalled in the previous section. In this case, the characteristic homomorphism c is not surjective over $\mathbb{Z}$ and γ_k $(1 \le k \le n)$ is not contained in the image of c. (Strictly speaking, we should consider the spinor group, because the special orthogonal group is not simply connected. In that case, γ_1 is contained in the image of c.) However, $2\gamma_k = c_k$ is contained in the image of c. So we can apply Theorem 3.2 to the polynomial $c_k = e_k(t_1, \ldots, t_n)$. Thus

$$c_k = \sum_{l(w)=k} \Delta_w(c_k) Z_w.$$

Let us compute $\Delta_w(c_k)$, where $l(w) = k$ for fixed k $(1 \le k \le n)$. For convenience of computation, we introduce the notation

$$c_l^{(m)} = e_l(t_1, t_2, \ldots, t_m) \quad (1 \le m \le n,\ 1 \le l \le m),$$

so that $c_k = c_k^{(n)}$.

We need the following auxiliary result.

Lemma 4.2. *For fixed k, $1 \le k \le n$, we have*

(1) $\Delta_i(c_k^{(n)}) = 0 \quad (1 \le i \le n-1)$,
(2) $\Delta_n(c_k^{(n)}) = 2c_{k-1}^{(n-1)}$,
(3) $\Delta_i(c_{k-j}^{(n-j)}) = 0 \quad (1 \le i \le n-j-1,\ 1 \le j \le n-1)$,
(4) $\Delta_{n-j}(c_{k-j}^{(n-j)}) = c_{k-j-1}^{(n-j-1)} \quad (1 \le j \le n-1)$.

Proof. (1) and (3) follow from the definition of Δ_i and the fact that $c_k^{(n)} = e_k(t_1, \ldots, t_n)$ (resp. $c_{k-j}^{(n-j)} = e_{k-j}(t_1, \ldots, t_{n-j})$) is invariant under the action of s_i $(1 \le i \le n-1)$ (resp. s_i $(1 \le i \le n-j-1)$).

By (4.3), we have, for $1 \le i \le n-1$,

$$\Delta_i(t_j) = \begin{cases} 1 & (j = i), \\ -1 & (j = i+1), \\ 0 & (j \ne i, i+1) \end{cases} \tag{4.5}$$

and

$$\Delta_n(t_j) = \begin{cases} 2 & (j = n), \\ 0 & (j \ne n). \end{cases} \tag{4.6}$$

Then we compute

$$\begin{aligned}\sum_{k=0}^{n}\Delta_n(c_k^{(n)}) &= \Delta_n\left(\prod_{i=1}^{n}(1+t_i)\right)\\ &= \Delta_n\left(\prod_{i=1}^{n-1}(1+t_i)\right)(1+t_n)+s_n\left(\prod_{i=1}^{n-1}(1+t_i)\right)\Delta_n(1+t_n)\\ &= 2\prod_{i=1}^{n-1}(1+t_i)\\ &= 2\sum_{k=0}^{n-1}c_k^{(n-1)}\end{aligned}$$

by (3.2) and (4.6). From this, (2) follows. (4) follows from a similar computation. □

By this lemma, we deduce that

$$\Delta_w(c_k) = \begin{cases} 2 & \text{if} \quad w = s_{n-k+1}\cdots s_{n-1}s_n, \\ 0 & \text{otherwise.}\end{cases}$$

Therefore, for $1 \leq k \leq n$, we have

$$c_k = 2Z_{n-k+1,\ldots,n-1,n}$$

in $H^*(SO(2n+1)/T^n;\mathbb{Z})$. Since γ_k is defined by

$$c_k = 2\gamma_k$$

and $H^*(SO(2n+1)/T^n;\mathbb{Z})$ is torsion free, we see that

$$\gamma_k = Z_{n-k+1,\ldots,n-1,n}.$$

Consequently, we obtain the following result.

Proposition 4.3. *In Theorem 4.1, the relation between the ring generators and the Schubert classes is given by*

$$\begin{aligned} t_1 &= Z_1,\\ t_i &= -Z_{i-1}+Z_i \quad (2\leq i\leq n-1),\\ t_n &= -Z_{n-1}+2Z_n,\\ \gamma_k &= Z_{n-k+1,\ldots,n-1,n} \quad (1\leq k\leq n).\end{aligned}$$

In particular, we can take $Z_1, Z_2, \ldots, Z_n, Z_{n-1,n}, \ldots, Z_{12\cdots n-1,n}$ *as the ring generators of* $H^*(SO(2n+1)/T^n;\mathbb{Z})$.

Remark 4.4. *The standard projection p from $SO(2n+1)/T^n$ to $SO(2n+1)/U(n)$ induces an injection*

$$p^*: H^*(SO(2n+1)/U(n);\mathbb{Z}) \hookrightarrow H^*(SO(2n+1)/T^n;\mathbb{Z}).$$

The cohomology ring $H^*(SO(2n+1)/U(n);\mathbb{Z})$ of the odd orthogonal Grassmannian $SO(2n+1)/U(n)$ has a $\mathbb{Z}$-basis of Schubert classes $\{\sigma_\lambda\}$ indexed by strict partitions λ contained in the "staircase" $\rho_n = (n, n-1, \ldots, 1)$. Observe that the generators $Z_{n-k+1,\ldots,n-1,n}$ $(1 \le k \le n)$ in Proposition 4.3 are the p^*-images of the "special Schubert classes" $\sigma_{(k)}$ $(1 \le k \le n)$ that were used by P. Pragacz to describe the cohomology ring of $SO(2n+1)/U(n)$ (see [17], *Theorem* 6.17*).*

The case of the even special orthogonal group $SO(2n)$ is almost identical to that of $SO(2n+1)$. So we only exhibit the data and results. Let

$$T^n = \left\{ \begin{pmatrix} \cos 2\pi t_1 & -\sin 2\pi t_1 & & & \\ \sin 2\pi t_1 & \cos 2\pi t_1 & & & \\ & & \ddots & & \\ & & & \cos 2\pi t_n & -\sin 2\pi t_n \\ & & & \sin 2\pi t_n & \cos 2\pi t_n \end{pmatrix} \right\}$$

be the standard maximal torus of $SO(2n)$. Then we have an isomorphism:

$$H^*(BT^n;\mathbb{Z}) = \mathbb{Z}[t_1, t_2, \ldots, t_n].$$

The system of simple roots is given by

$$\Pi = \{\alpha_1 = t_1 - t_2, \alpha_2 = t_2 - t_3, \ldots, \alpha_{n-1} = t_{n-1} - t_n, \alpha_n = t_{n-1} + t_n\}.$$

The corresponding fundamental weights $\{\omega_i\}_{1 \le i \le n}$ are given by

$$\begin{aligned} \omega_i &= t_1 + t_2 + \cdots + t_i \quad (1 \le i \le n-2), \\ \omega_{n-1} &= \frac{1}{2}(t_1 + \cdots + t_{n-1} - t_n), \\ \omega_n &= \frac{1}{2}(t_1 + \cdots + t_{n-1} + t_n). \end{aligned} \tag{4.7}$$

Let $s_i\,(1 \le i \le n)$ be the reflection corresponding to the simple root $\alpha_i\,(1 \le i \le n)$. Then the Weyl group $W(SO(2n))$ is finite and is generated by $s_i\,(1 \le i \le n)$ which act on $\{t_i\}_{1 \le i \le n}$ as permutations and an even number of sign changes:

$$\begin{aligned} W(SO(2n)) &= \langle s_1, s_2, \ldots, s_n \rangle \\ &\cong \mathcal{S}_n \ltimes (\mathbb{Z}/2\mathbb{Z})^{n-1}. \end{aligned} \tag{4.8}$$

The Borel presentation of $H^*(SO(2n)/T^n;\mathbb{Z})$ is given by

Theorem 4.5 (Toda-Watanabe [19], Corollary 2.2)**.** *The integral cohomology ring of $SO(2n)/T^n$ is*

$$H^*(SO(2n)/T^n;\mathbb{Z}) = \mathbb{Z}[t_1, t_2, \ldots, t_n, \gamma_1, \gamma_2, \ldots, \gamma_{n-1}] \Big/ \left(\begin{array}{l} c_i - 2\gamma_i \ (1 \le i \le n-1),\ c_n \\ \gamma_{2k} + \sum_{i=1}^{2k-1} (-1)^i \gamma_i \gamma_{2k-i} \ (1 \le k \le n-1) \end{array} \right),$$

where we denote by the same symbols $t_i \in H^2(SO(2n)/T^n;\mathbb{Z})$ the images of $t_i \in H^2(BT^n;\mathbb{Z})$ under the homomorphism c, $c_i = e_i(t_1, \ldots, t_n)$ $(1 \le i \le n)$, and $\gamma_i = 0$ for $i \ge n$.

By (4.7), we have

$$(4.9)\qquad \begin{aligned} t_1 &= \omega_1, \\ t_i &= -\omega_{i-1} + \omega_i \quad (2 \le i \le n-2), \\ t_{n-1} &= -\omega_{n-2} + \omega_{n-1} + \omega_n, \\ t_n &= -\omega_{n-1} + \omega_n. \end{aligned}$$

Since $c(\omega_i) = Z_i$, it follows from (4.9) that

$$(4.10)\qquad \begin{aligned} t_1 &= Z_1, \\ t_i &= -Z_{i-1} + Z_i \quad (2 \le i \le n-2), \\ t_{n-1} &= -Z_{n-2} + Z_{n-1} + Z_n, \\ t_n &= -Z_{n-1} + Z_n. \end{aligned}$$

By (4.9), we have, for $1 \le i \le n-1$,

$$(4.11)\qquad \Delta_i(t_j) = \begin{cases} 1 & (j = i), \\ -1 & (j = i+1), \\ 0 & (j \ne i, i+1) \end{cases}$$

and

$$(4.12)\qquad \Delta_n(t_j) = \begin{cases} 1 & (j = n-1), \\ 1 & (j = n), \\ 0 & (j \ne n-1, n). \end{cases}$$

Then we obtain the following quite similarly to Lemma 4.2.

Lemma 4.6. *For fixed k, $1 \le k \le n-1$, we have*

(1) $\Delta_i(c_k^{(n)}) = 0 \quad (1 \le i \le n-1)$,
(2) $\Delta_n(c_k^{(n)}) = 2c_{k-1}^{(n-2)}$,
(3) $\Delta_i(c_{k-j+1}^{(n-j)}) = 0 \quad (1 \le i \le n-j-1,\ 2 \le j \le n-1)$,

(4) $\Delta_{n-j}(c_{k-j+1}^{(n-j)}) = c_{k-j}^{(n-j-1)} \quad (2 \leq j \leq n-1)$.

By this lemma, we deduce that

$$\Delta_w(c_1) = \begin{cases} 2 & \text{if} \quad w = s_n, \\ 0 & \text{otherwise}; \end{cases}$$

$$\Delta_w(c_k) = \begin{cases} 2 & \text{if} \quad w = s_{n-k} \cdots s_{n-2}s_n, \\ 0 & \text{otherwise} \end{cases}$$

for $2 \leq k \leq n-1$.

Therefore we have

$$\begin{aligned} c_1 &= 2Z_n, \\ c_k &= 2Z_{n-k,\ldots,n-2,n} \quad (2 \leq k \leq n-1) \end{aligned}$$

in $H^*(SO(2n)/T^n; \mathbb{Z})$. Since γ_k is defined by

$$c_k = 2\gamma_k$$

and $H^*(SO(2n)/T^n; \mathbb{Z})$ is torsion free, we see that

$$\begin{aligned} \gamma_1 &= Z_n, \\ \gamma_k &= Z_{n-k,\ldots,n-2,n} \quad (2 \leq k \leq n-1). \end{aligned}$$

Consequently, we obtain the following result.

Proposition 4.7. *In Theorem 4.5, the relation between the ring generators and the Schubert classes is given by*

$$\begin{aligned} t_1 &= Z_1, \\ t_i &= -Z_{i-1} + Z_i \quad (2 \leq i \leq n-2), \\ t_{n-1} &= -Z_{n-2} + Z_{n-1} + Z_n, \\ t_n &= -Z_{n-1} + Z_n, \\ \gamma_1 &= Z_n, \\ \gamma_k &= Z_{n-k,\ldots,n-2,n} \quad (2 \leq k \leq n-1). \end{aligned}$$

In particular, we can take $Z_1, Z_2, \ldots, Z_n, Z_{n-2,n}, \ldots, Z_{12\cdots n-2,n}$ *as the ring generators of* $H^*(SO(2n)/T^n; \mathbb{Z})$.

Remark 4.8. *The standard projection* p *from* $SO(2n)/T^n$ *to* $SO(2n)/U(n)$ *induces an injection*

$$p^* : H^*(SO(2n)/U(n); \mathbb{Z}) \hookrightarrow H^*(SO(2n)/T^n; \mathbb{Z}).$$

The cohomology ring $H^*(SO(2n)/U(n); \mathbb{Z})$ *of the even orthogonal Grassmannian* $SO(2n)/U(n)$ *has a* $\mathbb{Z}$*-basis of Schubert classes* $\{\sigma_\lambda\}$ *indexed by strict partitions* λ *contained in* $\rho_{n-1} = (n-1, \ldots, 1)$. *Observe that the generators* $Z_n, Z_{n-k,\ldots,n-2,n}$ $(2 \leq k \leq n-1)$ *in Proposition 4.7* *are the* p^**-images of the special Schubert classes* $\sigma_{(k)}$ $(1 \leq k \leq n-1)$ *that were also used by P. Pragacz to describe the cohomology ring of* $SO(2n)/U(n)$ (*see* [17], *Theorem* 6.17′).

5. THE CASE OF G_2

In this section, we concentrate on the case of the exceptional Lie group G_2. Let T be a maximal torus of G_2. Following [4], we take the system of simple roots $\Pi = \{\alpha_1, \alpha_2\}$ and the corresponding fundamental weights $\{\omega_1, \omega_2\}$. Then we can identify

$$H^*(BT;\mathbb{Z}) = \mathbb{Z}[\omega_1, \omega_2].$$

Let $s_i\,(i = 1, 2)$ be the reflection corresponding to the simple root $\alpha_i\,(i = 1, 2)$. Then the Weyl group $W(G_2)$ of G_2 is finite and is generated by $s_i\,(i = 1, 2)$:

$$\begin{aligned} &W(G_2) = \langle s_1, s_2 \rangle, \\ &s_1^2 = s_2^2 = 1,\ (s_1 s_2)^6 = 1. \end{aligned} \tag{5.1}$$

Now we review the Borel presentation of $H^*(G_2/T;\mathbb{Z})$. We put

$$\begin{aligned} t_1 &= -\omega_1, \\ t_2 &= -\omega_1 + \omega_2, \\ t_3 &= 2\omega_1 - \omega_2, \\ c_i &= e_i(t_1, t_2, t_3). \end{aligned} \tag{5.2}$$

Then we can write

$$H^*(BT;\mathbb{Z}) = \mathbb{Z}[t_1, t_2, t_3]/(c_1).$$

The action of $W(G_2)$ on $\{t_1, t_2, t_3\}$ is given by TABLE 1.

	s_1	s_2
t_1	$-t_2$	t_1
t_2	$-t_1$	t_3
t_3	$-t_3$	t_2

TABLE 1.

Remark 5.1. *The elements* $\{t_i\}_{i=1,2,3}$ *are derived from the natural inclusion* $T \subset SU(3) \subset G_2$.

The integral cohomology ring of G_2/T was first determined by Bott-Samelson [3], but we prefer to use the presentation due to Toda-Watanabe.

Theorem 5.2 (Bott-Samelson [3], Toda-Watanabe [19])**.** *The integral cohomology ring of* G_2/T *is*

$$H^*(G_2/T;\mathbb{Z}) = \mathbb{Z}[t_1, t_2, t_3, \gamma_3]/(\rho_1, \rho_2, \rho_3, \rho_6),$$

where $\rho_1 = c_1,\ \rho_2 = c_2,\ \rho_3 = c_3 - 2\gamma_3,\ \rho_6 = \gamma_3^2$, *and we denote by the same symbols* $t_i \in H^2(G_2/T;\mathbb{Z})$ *the images of* $t_i \in H^2(BT;\mathbb{Z})$ *under the homomorphism* c.

By (5.1), the elements of the Weyl group $W(G_2)$ are given by the following table.

$l(w)$	Elements of $W(G_2)$	
0	1	
1	s_1	s_2
2	s_1s_2	s_2s_1
3	$s_1s_2s_1$	$s_2s_1s_2$
4	$s_1s_2s_1s_2$	$s_2s_1s_2s_1$
5	$s_1s_2s_1s_2s_1$	$s_2s_1s_2s_1s_2$
6	$s_1s_2s_1s_2s_1s_2$	

Therefore the corresponding Schubert basis for $H^*(G_2/T;\mathbb{Z})$ is given as follows:

deg	0	2	4	6	8	10	12
	1	Z_1	Z_{12}	Z_{121}	Z_{1212}	Z_{12121}	
		Z_2	Z_{21}	Z_{212}	Z_{2121}	Z_{21212}	Z_{121212}

Here we denote Z_{s_i} simply by Z_i and so on. We wish to express the algebra generators $\{t_1, t_2, t_3, \gamma_3\}$ in terms of Schubert classes. Since $c(\omega_i) = Z_i$ $(i = 1, 2)$, it follows from (5.2) that

$$\begin{aligned} t_1 &= -Z_1, \\ t_2 &= -Z_1 + Z_2, \\ t_3 &= 2Z_1 - Z_2. \end{aligned} \tag{5.3}$$

Next we can put

$$\gamma_3 = a_{121}Z_{121} + a_{212}Z_{212}$$

for some integers a_{121}, a_{212} and we need to determine the coefficients a_{121}, a_{212}. The characteristic homomorphism c is not surjective over $\mathbb{Z}$ and γ_3 is not contained in the image of c, but $2\gamma_3 = c_3$ is in the image of c. Thus putting

$$c_3 = b_{121}Z_{121} + b_{212}Z_{212}$$

for some integers b_{121}, b_{212}, we can compute the coefficients b_{121}, b_{212} using the divided difference operators. By (5.2), we have

$$\begin{aligned} c_3 &= t_1t_2t_3 \\ &= 2\omega_1^3 - 3\omega_1^2\omega_2 + \omega_1\omega_2^2. \end{aligned}$$

Therefore we derive

$$\begin{aligned} b_{121} &= \Delta_1\Delta_2\Delta_1(c_3) \\ &= \Delta_1\Delta_2\Delta_1(2\omega_1^3 - 3\omega_1^2\omega_2 + \omega_1\omega_2^2) \\ &= -2, \\ b_{212} &= \Delta_2\Delta_1\Delta_2(c_3) \\ &= \Delta_2\Delta_1\Delta_2(2\omega_1^3 - 3\omega_1^2\omega_2 + \omega_1\omega_2^2) \\ &= 0 \end{aligned}$$

from (3.1), (3.2) and Table 1.

Thus we have

$$c_3 = -2Z_{121}$$

in $H^6(G_2/T;\mathbb{Z})$. Since γ_3 is defined by

$$c_3 = 2\gamma_3$$

and $H^*(G_2/T;\mathbb{Z})$ is torsion free, we see that

$$\gamma_3 = -Z_{121}.$$

Consequently, we obtain the following result.

Proposition 5.3. *In Theorem 5.2, the relation between the ring generators* $\{t_1, t_2, t_3, \gamma_3\}$ *and the Schubert classes is given by*

$$\begin{aligned} t_1 &= -Z_1, \\ t_2 &= -Z_1 + Z_2, \\ t_3 &= 2Z_1 - Z_2, \\ \gamma_3 &= -Z_{121}. \end{aligned}$$

In particular, we can take Z_1, Z_2, Z_{121} *as the ring generators of* $H^*(G_2/T;\mathbb{Z})$.

6. The case of F_4

In this section, we deal with the case of the exceptional Lie group F_4. Let T be a maximal torus of F_4. Following [4], we take the system of simple roots $\Pi = \{\alpha_i\}_{1\le i\le 4}$ and the corresponding fundamental weights $\{\omega_i\}_{1\le i\le 4}$. Then we can identify

$$H^*(BT;\mathbb{Z}) = \mathbb{Z}[\omega_1, \omega_2, \omega_3, \omega_4].$$

Let $s_i\,(1 \leq i \leq 4)$ be the reflection corresponding to the simple root $\alpha_i\,(1 \leq i \leq 4)$. Then the Weyl group $W(F_4)$ of F_4 is finite and is generated by $s_i\,(1 \leq i \leq 4)$:

$$\begin{aligned} &W(F_4) = \langle s_1, s_2, s_3, s_4 \rangle, \\ &s_1^2 = s_2^2 = s_3^2 = s_4^2 = 1, \\ &(s_1 s_2)^3 = (s_3 s_4)^3 = (s_2 s_3)^4 = 1, \\ &s_1 s_3 = s_3 s_1, s_1 s_4 = s_4 s_1, s_2 s_4 = s_4 s_2. \end{aligned} \tag{6.1}$$

Now we review the Borel presentation of $H^*(F_4/T;\mathbb{Z})$. We put

$$\begin{aligned} t_1 &= -\omega_4, \\ t_2 &= \omega_1 - \omega_4, \\ t_3 &= -\omega_1 + \omega_2 - \omega_4, \\ t_4 &= -\omega_2 + 2\omega_3 - \omega_4, \\ c_i &= e_i(t_1, \ldots, t_4), \\ t &= \frac{1}{2}c_1 = \omega_3 - 2\omega_4. \end{aligned} \tag{6.2}$$

Then we can write

$$H^*(BT;\mathbb{Z}) = \mathbb{Z}[t_1, t_2, t_3, t_4, t]/(c_1 - 2t).$$

The action of $W(F_4)$ on $\{t_i\}_{1 \leq i \leq 4}$ is given by Table 2, where blanks indicate the trivial action.

	s_1	s_2	s_3	s_4
t_1				$t_1 - t$
t_2	t_3			$t_2 - t$
t_3	t_2	t_4		$t_3 - t$
t_4		t_3	$-t_4$	$t_4 - t$
t			$t - t_4$	$-t$

Table 2.

Remark 6.1. *The elements $\{t_i\}_{1 \leq i \leq 4}$ and t are derived from the natural inclusion $T \subset Spin(9) \subset F_4$.*

The integral cohomology ring of F_4/T was determined by Toda-Watanabe [19].

Theorem 6.2 (Toda-Watanabe [19], Theorem A)**.** *The integral cohomology ring of F_4/T is*

$$H^*(F_4/T;\mathbb{Z}) = \mathbb{Z}[t_1, t_2, t_3, t_4, t, \gamma_3, \gamma_4]/(\rho_1, \rho_2, \rho_3, \rho_4, \rho_6, \rho_8, \rho_{12}),$$

where

$$\begin{aligned}
\rho_1 &= c_1 - 2t, \\
\rho_2 &= c_2 - 2t^2, \\
\rho_3 &= c_3 - 2\gamma_3, \\
\rho_4 &= c_4 - 4t\gamma_3 + 8t^4 - 3\gamma_4, \\
\rho_6 &= \gamma_3^2 - 3t^2\gamma_4 - 4t^3\gamma_3 + 8t^6, \\
\rho_8 &= 3\gamma_4^2 + 6t\gamma_3\gamma_4 - 3t^4\gamma_4 - 13t^8, \\
\rho_{12} &= \gamma_4^3 - 6t^4\gamma_4^2 + 12t^8\gamma_4 - 8t^{12},
\end{aligned}$$

and we denote by the same symbols $t_i \in H^2(F_4/T;\mathbb{Z})$ *the images of* $t_i \in H^2(BT;\mathbb{Z})$ *under the homomorphism* c*.*

Remark 6.3. *In* [19]*, Toda and Watanabe described* $H^*(F_4/T;\mathbb{Z})$ *using the inclusion* $H^*(F_4/Spin(9);\mathbb{Z}) \hookrightarrow H^*(F_4/T;\mathbb{Z})$ *and the known structure of* $H^*(F_4/Spin(9);\mathbb{Z})$*. Theorem 6.2 is a rewritten form of their result in terms of the* $W(F_4)$*-invariants.*

By (6.1), the elements of the Weyl group $W(F_4)$ of length ≤ 4 are given by the following table.

$l(w)$	Elements of $W(F_4)$				
0	1				
1	s_1	s_2	s_3	s_4	
2	s_1s_2	s_1s_3	s_1s_4	s_2s_1	s_2s_3
	s_2s_4	s_3s_2	s_3s_4	s_4s_3	
3	$s_1s_2s_1$	$s_1s_2s_3$	$s_1s_2s_4$	$s_1s_3s_2$	$s_1s_3s_4$
	$s_1s_4s_3$	$s_2s_1s_3$	$s_2s_1s_4$	$s_2s_3s_2$	$s_2s_3s_4$
	$s_2s_4s_3$	$s_3s_2s_1$	$s_3s_2s_3$	$s_3s_2s_4$	$s_3s_4s_3$
	$s_4s_3s_2$				
4	$s_1s_2s_1s_3$	$s_1s_2s_1s_4$	$s_1s_2s_3s_2$	$s_1s_2s_3s_4$	$s_1s_2s_4s_3$
	$s_1s_3s_2s_1$	$s_1s_3s_2s_3$	$s_1s_3s_2s_4$	$s_1s_3s_4s_3$	$s_1s_4s_3s_2$
	$s_2s_1s_3s_2$	$s_2s_1s_3s_4$	$s_2s_1s_4s_3$	$s_2s_3s_2s_1$	$s_2s_3s_2s_3$
	$s_2s_3s_2s_4$	$s_2s_3s_4s_3$	$s_2s_4s_3s_2$	$s_3s_2s_1s_3$	$s_3s_2s_1s_4$
	$s_3s_2s_3s_4$	$s_3s_2s_4s_3$	$s_3s_4s_3s_2$	$s_4s_3s_2s_1$	$s_4s_3s_2s_3$

We have the corresponding Schubert basis $\{Z_w\}_{w\in W(F_4)}$. As before, we denote Z_{s_i} simply by Z_i and so on. We wish to express the algebra generators $\{t_1, t_2, t_3, t_4, t, \gamma_3, \gamma_4\}$ in terms of Schubert classes. Since $c(\omega_i) = Z_i$, it follows from (6.2) that

$$
\begin{aligned}
t_1 &= -Z_4, \\
t_2 &= Z_1 - Z_4, \\
t_3 &= -Z_1 + Z_2 - Z_4, \\
t_4 &= -Z_2 + 2Z_3 - Z_4, \\
t &= Z_3 - 2Z_4.
\end{aligned} \tag{6.3}
$$

Next we can put

$$\gamma_3 = \sum_{l(w)=3} a_w Z_w, \quad \gamma_4 = \sum_{l(w)=4} a_w Z_w$$

for some integers a_w. We wish to determine the coefficients a_w. By Theorem 6.2, we have

$$
\begin{aligned}
2\gamma_3 &= c_3, \\
3\gamma_4 &= c_4 - 4t\gamma_3 + 8t^4 \\
&= c_4 - 2tc_3 + 8t^4.
\end{aligned} \tag{6.4}
$$

Therefore $2\gamma_3$ and $3\gamma_4$ are contained in the image of c. So as in the case of G_2, we apply the divided difference operators to the right hand side of (6.4). The result can be seen in the following table.

	c_3
Δ_{121}	0
Δ_{123}	2
Δ_{124}	0
Δ_{132}	0
Δ_{134}	0
Δ_{143}	0
Δ_{213}	0
Δ_{214}	0
Δ_{232}	0
Δ_{234}	-2
Δ_{243}	-4
Δ_{321}	0
Δ_{323}	0
Δ_{324}	0
Δ_{343}	6
Δ_{432}	0

Here we denote $\Delta_1\Delta_2\Delta_1$ simply by Δ_{121} and so on.

Thus we have

$$\begin{aligned} c_3 &= 2Z_{123} - 2Z_{234} - 4Z_{243} + 6Z_{343} \\ &= 2(Z_{123} - Z_{234} - 2Z_{243} + 3Z_{343}) \end{aligned}$$

in $H^6(F_4/T;\mathbb{Z})$. Since γ_3 is defined by

$$c_3 = 2\gamma_3$$

and $H^*(F_4/T;\mathbb{Z})$ is torsion free, we see that

$$\gamma_3 = Z_{123} - Z_{234} - 2Z_{243} + 3Z_{343}. \tag{6.5}$$

Similarly we obtain

	$c_4 - 2tc_3 + 8t^4$
Δ_{1213}	0
Δ_{1214}	0
Δ_{1232}	0
Δ_{1234}	3
Δ_{1243}	-30
Δ_{1321}	0
Δ_{1323}	12
Δ_{1324}	0
Δ_{1343}	0
Δ_{1432}	0
Δ_{2132}	0
Δ_{2134}	0
Δ_{2143}	0
Δ_{2321}	0
Δ_{2323}	0
Δ_{2324}	0
Δ_{2343}	0
Δ_{2432}	0
Δ_{3213}	0
Δ_{3214}	0
Δ_{3234}	-3
Δ_{3243}	30
Δ_{3432}	0
Δ_{4321}	0
Δ_{4323}	-24

Thus we have

$$\begin{aligned} c_4 - 2tc_3 + 8t^4 &= 3Z_{1234} - 30Z_{1243} + 12Z_{1323} - 3Z_{3234} + 30Z_{3243} - 24Z_{4323} \\ &= 3(Z_{1234} - 10Z_{1243} + 4Z_{1323} - Z_{3234} + 10Z_{3243} - 8Z_{4323}) \end{aligned}$$

in $H^8(F_4/T;\mathbb{Z})$. Since γ_4 is defined by

$$c_4 - 2tc_3 + 8t^4 = 3\gamma_4$$

and $H^*(F_4/T;\mathbb{Z})$ is torsion free, we see that

$$\gamma_4 = Z_{1234} - 10Z_{1243} + 4Z_{1323} - Z_{3234} + 10Z_{3243} - 8Z_{4323}. \tag{6.6}$$

Unfortunately, from (6.5) and (6.6), we cannot decide which Schubert classes are indecomposable. This leads us to the use of the Giambelli formula. We need the following data.

(1) A set of positive roots of F_4 is given by

$$\Delta^+ = \left\{ \begin{array}{l} t_i \pm t_j \ (1 \leq i < j \leq 4), \\ t_i \ (1 \leq i \leq 4), \ \dfrac{1}{2}(t_1 \pm t_2 \pm t_3 \pm t_4) \end{array} \right\}.$$

(2) The longest element w_0 of the Weyl group $W(F_4)$ is given by

$$w_0 = s_1s_2s_1s_3s_2s_1s_3s_2s_3s_4s_3s_2s_1s_3s_2s_3s_4s_3s_2s_1s_3s_2s_3s_4.$$

Then, using the Giambelli formula (Theorem 3.3), we obtain the following result.

Lemma 6.4. *In (6.5) and (6.6), each Schubert class is expressed in terms of* t_i $(1 \leq i \leq 4), t, \gamma_3, \gamma_4$ *as follows*:

$$\begin{aligned} Z_{123} &= \gamma_3 - 2t_1^3 + 3t_1^2t - 2t_1t^2, \\ Z_{234} &= -t_1^3, \\ Z_{243} &= -2t_1^3 + 3t_1^2t - t_1t^2, \\ Z_{343} &= -t_1^3 + t_1^2t, \\ Z_{1234} &= -\gamma_4 + (t_1 - 2t)\gamma_3 + t_1^3t - t_1^2t^2 + 3t^4, \\ Z_{1243} &= \gamma_4 + (-2t_1 + 2t)\gamma_3 + 2t_1^4 - 4t_1^3t + 3t_1^2t^2 - 3t^4, \\ Z_{1323} &= -\gamma_4 - t\gamma_3 + 2t_1^4 - 4t_1^3t + 4t_1^2t^2 - 2t_1t^3 + 3t^4, \\ Z_{3234} &= \gamma_4 + (-t_1 + 2t)\gamma_3 + t_1^4 - t_1^3t + t_1^2t^2 - 3t^4, \\ Z_{3243} &= t_1^4 - 2t_1^3t + t_1^2t^2, \\ Z_{4323} &= -\gamma_4 + (2t_1 - 2t)\gamma_3 - t_1t^3 + 3t^4. \end{aligned}$$

In particular, Z_{123} *and* Z_{1234} *are indecomposable in the ring* $H^*(F_4/T;\mathbb{Z})$*.*

Consequently, we obtain the following result.

Proposition 6.5. *In Theorem 6.2, the relation between the ring generators* $\{t_1, t_2, t_3, t_4, t, \gamma_3, \gamma_4\}$ *and the Schubert classes is given by*

$$\begin{aligned}
t_1 &= -Z_4,\\
t_2 &= Z_1 - Z_4,\\
t_3 &= -Z_1 + Z_2 - Z_4,\\
t_4 &= -Z_2 + 2Z_3 - Z_4,\\
t &= Z_3 - 2Z_4,\\
\gamma_3 &= Z_{123} - Z_{234} - 2Z_{243} + 3Z_{343},\\
\gamma_4 &= Z_{1234} - 10Z_{1243} + 4Z_{1323} - Z_{3234} + 10Z_{3243} - 8Z_{4323}.
\end{aligned}$$

Furthermore, we have

$$\begin{aligned}
Z_{123} &= \gamma_3 - 2t_1^3 + 3t_1^2 t - 2t_1 t^2,\\
Z_{1234} &= -\gamma_4 + (t_1 - 2t)\gamma_3 + t_1^3 t - t_1^2 t^2 + 3t^4,
\end{aligned}$$

and we can take $Z_1, Z_2, Z_3, Z_4, Z_{123}, Z_{1234}$ *as the ring generators of* $H^*(F_4/T;\mathbb{Z})$.

7. The Chow rings of $\mathrm{SO}(n), \mathrm{Spin}(n), \mathrm{G}_2$ and F_4

In this section, using our description of the integral cohomology rings of the flag manifolds of types B_n, D_n, G_2 and F_4 (see Sections 4, 5 and 6) and a remark of Grothendieck, we compute the Chow rings of the corresponding complex algebraic groups.

As in Section 2, let K be a compact simply connected simple Lie group, T its maximal torus, $G = K^{\mathbb{C}}$ the complexification of K and B a Borel subgroup of G containing T. Let $A^i(\cdot)$ denote the Chow group of codimension i cycles up to rational equivalence and $A(\cdot) = \bigoplus_{i \geq 0} A^i(\cdot)$. Given any character χ of B, that is, a homomorphism of B into $\mathbb{C}^\times$, we have the associated line bundle L_χ over G/B, which defines an element of $A^1(G/B)$, denoted by $c(\chi)$. This induces a homomorphism

$$c : \hat{B} \longrightarrow A^1(G/B),$$

where $\hat{B}$ denotes the character group of B. Extending this homomorphism c by multiplicativity to the symmetric algebra $S(\hat{B})$ of $\hat{B}$, one obtains a homomorphism

$$c : S(\hat{B}) \longrightarrow A(G/B), \tag{7.1}$$

which is also called the *characteristic homomorphism.*

Then Grothendieck's remark ([9], p.21, Remarques 2°) allows us to obtain $A(G)$ as the quotient of $A(G/B)$ by the ideal generated by $c(\hat{B})$. Denote by $T_G : A(G/B) \longrightarrow A(G)$ the canonical map onto the quotient. It is known ([9], Lemme 10) that the Chow ring $A(G/B)$ of G/B is isomorphic to the integral cohomology ring $H^*(G/B;\mathbb{Z})$ of G/B. Under this isomorphism, the Schubert variety X_{w_0w} corresponds to the Schubert class Z_w and the above characteristic homomorphism (7.1) coincides with the characteristic homomorphism (2.1).

Thus, in order to determine the Chow ring $A(G)$, we need only to compute the quotient ring of $H^*(K/T;\mathbb{Z})$ by the ideal generated by $c(H^2(BT;\mathbb{Z}))$. Since we assume that K is simply connected, $H^2(BT;\mathbb{Z})$ is isomorphic to $H^2(K/T;\mathbb{Z})$. Therefore we compute the quotient ring $H^*(K/T;\mathbb{Z})/(H^2(K/T;\mathbb{Z}))$. We will show how to do this for F_4. By Theorem 6.2, we have

$$H^*(F_4/T;\mathbb{Z})/(t_1,t_2,t_3,t_4,t) \quad = \mathbb{Z}[\gamma_3,\gamma_4]/(2\gamma_3, 3\gamma_4, \gamma_3^2, \gamma_4^3).$$

Taking Proposition 6.5 into account, we can replace γ_3, γ_4 with Z_{123}, Z_{1234} respectively. Thus we obtain the following

Theorem 7.1. *If G is of type* F_4*, we have*

$$A(\mathrm{F}_4) = \mathbb{Z}[X_3, X_4]/(2X_3, 3X_4, X_3^2, X_4^3),$$

where X_3 (resp. X_4) is the image under T_G of the element of $A(G/B)$ defined by the Schubert variety $X_{w_0s_1s_2s_3}$ (resp. $X_{w_0s_1s_2s_3s_4}$).

In a similar way, we can compute the Chow rings of $\mathrm{SO}(n), \mathrm{Spin}(n)$ and G_2. The results are summarized as follows.

Theorem 7.2. *If G is of type B_n $(\mathrm{SO}(2n+1)$ or $\mathrm{Spin}(2n+1))$ and x_i $(1 \leq i \leq n)$ is the element of $A(G/B)$ defined by the Schubert variety $X_{w_0s_{n-i+1}\cdots s_{n-1}s_n}$. Then we have*

$$A(\mathrm{SO}(2n+1)) = \mathbb{Z}[X_1, X_3, X_5, \ldots, X_{2[\frac{n+1}{2}]-1}]/(2X_i,\ X_i^{p_i}),$$
$$A(\mathrm{Spin}(2n+1)) = \mathbb{Z}[X_3, X_5, \ldots, X_{2[\frac{n+1}{2}]-1}]/(2X_i,\ X_i^{p_i}),$$

where

$$p_i = 2^{[\log_2 \frac{n}{i}]+1}, \quad X_i = T_G(x_i).$$

Theorem 7.3. *If G is of type D_n $(\mathrm{SO}(2n)$ or $\mathrm{Spin}(2n))$ and x_1 (resp. x_i $(2 \leq i \leq n-1)$) is the element of $A(G/B)$ defined by the Schubert variety $X_{w_0s_n}$ (resp. $X_{w_0s_{n-i}\cdots s_{n-2}s_n}$ $(2 \leq i \leq n-1)$). Then we have*

$$A(\mathrm{SO}(2n)) = \mathbb{Z}[X_1, X_3, X_5, \ldots, X_{2[\frac{n}{2}]-1}]/(2X_i,\ X_i^{p_i}),$$
$$A(\mathrm{Spin}(2n)) = \mathbb{Z}[X_3, X_5, \ldots, X_{2[\frac{n}{2}]-1}]/(2X_i,\ X_i^{p_i}),$$

where

$$p_i = 2^{[\log_2 \frac{n-1}{i}]+1}, \quad X_i = T_G(x_i).$$

Theorem 7.4. *If G is of type G_2, we have*

$$A(\mathrm{G}_2) = \mathbb{Z}[X_3]/(2X_3, X_3^2),$$

where X_3 is the image under T_G of the element of $A(G/B)$ defined by the Schubert variety $X_{w_0 s_1 s_2 s_1}$.

We observe that our results obtained in this section agree with those obtained in Marlin [13].

REFERENCES

[1] I.N. Bernstein, I.M. Gelfand and S.I. Gelfand, *Schubert cells, and the cohomology of the spaces G/P*, London Math. Soc. Lecture Note Ser. **69** Cambridge Univ. Press, 1982, 115-140.

[2] A. Borel, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Ann. of Math. **57** (1953), 115-207.

[3] R. Bott and H. Samelson, *The integral cohomology ring of G/T*, Proc. Nat. Acad. Sci. USA **41** (1955), 490-493.

[4] N. Bourbaki, *Groupes et Algèbre de Lie. Chapitre* 4, 5 *et* 6, Masson, Paris, 1981.

[5] C. Chevalley, Sur les décomposition cellulaires des espaces G/B, Algebraic Groups and their Generalizations: Classical Methods (W. Haboush, ed.), Proc. Sympos. Pure Math., **56**, Part 1, Amer. Math. Soc., 1994, 1-23.

[6] M. Demazure, *Invariants symétriques entiers des groupes de Weyl et torsion*, Invent. Math. **21** (1973), 287-301.

[7] H. Duan, *Multiplicative rule of Schubert classes*, Invent. Math. **159** (2005), 407-436.

[8] H. Duan and X. Zhao, *The Chow rings of generalized Grassmannians*, arXiv:math.AG/0511332.

[9] A. Grothendieck, *Torsion homologique et sections rationnelles*, Exposé 5 in Anneaux de Chow et applications, Séminaire C. Chevalley, 1958, 1-29, Multigraphié, Secrétariat mathématique, Paris.

[10] H. Hiller, *Schubert calculus of a Coxeter group*, Enseign. Math. (2) **27** (1981), 57-84.

[11] A. Iliev and L. Manivel, *The Chow ring of the Cayley planes*, Compos. Math. **141** (2005), 146-160.

[12] S. Kaji and M. Nakagawa *The Chow rings of the algebraic groups* E_6 *and* E_7, arXiv:math.AT/07093702.

[13] R. Marlin, *Anneaux de Chow des groupes algebriques* $\mathrm{SO}(n), \mathrm{Spin}(n), \mathrm{G}_2$ *et* F_4, Publications Math. d'Orsay, 95-7419 (1974).

[14] M. Nakagawa, *The integral cohomology ring of E_7/T*, J. Math. Kyoto Univ. **41** (2001), 303-321.

[15] M. Nakagawa, *The integral cohomology ring of E_8/T*, in preparation.

[16] P. Pragacz, *Multiplying Schubert classes*, Topics in Cohomological Studies of Algebraic Varieties, 163-174, Trends in Math., Birkhäuser Verlag Basel, 2005.

[17] P. Pragacz, *Algebro-geometric applications of Schur S- and Q-polynomials*, Topics in invariant theory (M.P. Malliavin, ed.), Lecture Notes in Math., 1478, 130-191, Springer Verlag, 1991.

[18] H. Toda, *On the cohomology ring of some homogeneous spaces*, J. Math. Kyoto Univ. **15** (1975), 185-199.

[19] H. Toda and T. Watanabe, *The integral cohomology ring of* F_4/T *and* E_6/T, J. Math. Kyoto Univ. **14** (1974), 257-286.

DEPARTMENT OF GENERAL EDUCATION
TAKAMATSU NATIONAL COLLEGE OF TECHNOLOGY
TAKAMATSU 761-8058, JAPAN
E-mail address: `nakagawa@takamatsu-nct.ac.jp`